



LOWER YOUR BLOOD PRESSURE NATURALLY

VITAL STEPS ANYONE CAN FOLLOW

Dr. James Meschino DC, MS, ND



- A Comprehensive Guide to Lowering Blood Pressure Naturally
- A Comprehensive Guide to Lowering Blood Pressure Naturally
- A Comprehensive Guide to Lowering Blood Pressure Naturally

About Dr. James Meschino, DC, MS, ND



A recognized expert in the use of nutritional supplements in the prevention and management of degenerative diseases and anti-aging, Dr. James Meschino, DC, MS, ND, was appointed to the advisory board of the Academy of Anti-Aging Research in 2001. He is a doctor of naturopathy, an associate professor at the Canadian Memorial Chiropractic College and has been a Faculty Member of the American Council of Exercise (ACE). He is also a faculty member of the Integrative Cancer Therapy Fellowship Program for physicians, sanctioned by the American Academy of Anti-Aging Medicine.

Dr. Meschino has appeared as a health and anti-aging expert on many television and radio programs in Canada and the United States.

The published author of five nutrition, supplementation and wellness books, he has also had over 50 research review papers on nutritional supplementation published by *America-Online* and is the regular anti-aging and natural therapies columnist for *Dynamic Chiropractic*. Dr. Meschino's continuing education seminars for health practitioners are authorized for continuing education credits in many states and provinces throughout North America.

- A Comprehensive Guide to Minerals
- A Comprehensive Guide to Herbs
- A Comprehensive Guide to Accessory Nutrients and Esser

Table of Contents

(move your mouse over text below, then click to follow link)

INTRODUCTION	page 4
TRENDS IN HYPERTENSION	page 4
NUTRITIONAL THERAPIES & LIFESTYLE INTERVENTIONS	pages 5 - 8
Lose Weight	page 5
Lower Alcohol Consumption	page 6
Sodium Restriction	page 6
Calcium Supplementation	page 6
Magnesium Supplementation	page 7
Omega-3 Fat Supplementation	page 7
Garlic Extract Supplementation	page 7
Coenzyme Q10 Supplementation	page 7
Hawthorn Extract Supplementation	page 8
SUMMARY	page 8
WORLD HEALTH ORGANIZATION RECOMMENDATIONS	page 9
ADDITIONAL READINGS	page 10
SCIENTIFIC REFERENCES	page 11

Introduction


In day-to-day practice, many practitioners encounter patients with hypertension problems that are not being managed effectively or where the patient has refused to continue their medications due to undesirable side effects or for other reasons. Natural health practitioners are often asked if there are any dietary supplements or nutritional therapies that can lower blood pressure in a more natural way without producing unwanted side effects.

Research studies conducted over the past fifteen years supports the use of specific dietary and supplementation practices, as well as participation in physical activity, as natural interventions to reduce high blood pressure. In some cases these natural solutions are all that are required to control blood pressure, and in other cases these practices can significantly lower the requirement for medication, helping to reduce the likelihood of adverse side effects occurring from the use of these drugs.

Research studies conducted over the past fifteen years supports the use of specific dietary and supplementation practices, as well as participation in physical activity, as natural interventions to reduce high blood pressure.

Trends in Hypertension

High blood pressure affects approximately twenty-five percent of the adult population in developed countries like the United States and Canada. In up to 75% of these cases hypertension manifests in a mild form, which is highly



sensitive to nutrition, supplementation and lifestyle practices.^(1,22) Even the most current medical literature stresses that all persons with documented hypertension should receive intensive non-pharmacologic therapies to improve control of their condition and reduce their risk of developing further cardiovascular disease.⁽²⁾

Hypertension, along with hypercholesterolemia and cigarette smoking are considered to be the three cardinal risk factors for cardiovascular disease. Studies indicate that lowering a patient's blood pressure from 160/90 to 140/80mmHg may decrease risk of heart disease by more than 30 percent.⁽³⁾

Studies indicate that lowering a patient's blood pressure from 160/90 to 140/80mmHg may decrease risk of heart disease by more than 30 percent.

From a medical standpoint, the use of anti-hypertensive drugs dominates the management of these conditions and little attention is often given to nutrition and lifestyle approaches. However, many patients discontinue their drug regime due to




side effects from these drugs, which can include fatigue, male impotence, elevated cholesterol levels, light-headedness, dizziness and skin eruptions. ⁽⁴⁾ In Canada, 22% of adults have hypertension, but only 16% of this population are treated and controlled. This leaves 84% of hypertensive patients uncontrolled and sometimes unaware that this silent killer is even present. ^(5,6) In general, hypertension

across the population is not well controlled and an effort by alternative health care providers to help remedy this situation is urgently needed as cardiovascular disease continues to be the leading cause of premature death in our society.



Effective Nutritional Therapies and Lifestyle Interventions

1. **Lose Weight** – Hypertensive patients who are overweight experience a drop to normal in their readings in approximately two-thirds of cases by simply losing 10-15 pounds. ^(7,8) Overweight patients tend to display insulin resistance, especially in cases where there is a propensity for abdominal weight gain (android obesity). Insulin resistance results in higher secretion rates of insulin displayed by peripheral body cells. One of the consequences of hyperinsulinemia is increased retention of sodium by the kidneys, which tends to drive up blood pressure in sodium-sensitive individuals. Thus, moderate weight loss helps to reverse insulin resistance, lowering basal and postprandial insulin blood levels. This, in turn, encourages less sodium retention and a natural lowering of blood pressure in these individuals. It is estimated that in up to 50% of adults in the United States whose hypertension is being pharmacologically managed, the need for drug therapy could be alleviated with only modest reductions in body weight. ⁽⁹⁾



Hypertensive patients who are overweight experience a drop to normal in their readings in approximately two-thirds of cases by simply losing 10-15 pounds.

In conjunction with dietary advice to help reduce excess weight, engaging in regular endurance-based exercise (optimum 40 to 60 minutes of at least brisk walking, four to five times per week) has been shown to help reduce high blood pressure. Exercise further increases insulin sensitivity, accelerates weight loss and induces other



changes within the cardiovascular system which lend themselves to a lowering of blood pressure^(6,10) Clearly, health practitioners should become more involved in providing patients with safe and effective nutrition and lifestyle practices that reverse weight gain and enhance the patient's overall level of cardiovascular fitness.

2. **Lower Alcohol Consumption** — Studies indicate that excess alcohol consumption is a culprit in hypertension. Restricting alcohol consumption to two or fewer drinks per day; fewer than 14 drinks weekly for men, 9 for women, has been shown to help lower blood pressure in individuals who consume alcohol.⁽⁷⁾



3. **Sodium Restriction** — Approximately 40-50% of hypertensive patients are thought to be sensitive to sodium intake as at least a partial cause of their problem. Salt sensitivity appears to be more common among blacks, diabetics and the elderly. Reducing sodium intake to 2000 mg per day is a prudent step in the global management of hypertension. This requires restricted use of discretionary salt and avoiding heavily salted processed foods. (e.g., prepared soups, pickles, salted snack foods, foods containing MSG etc.)^(7,11,12,13)

4. **Calcium Supplementation** — A number of well-designed human intervention trials reveal that calcium supplementation (1,000-1,500 mg calcium per day as calcium carbonate or citrate) can lower blood pressure, particularly in sodium-sensitive hypertensive patients. Calcium encourages sodium excretion by the kidneys and, in concert with magnesium, helps to relax the smooth muscle lining of arterioles, lowering diastolic pressure.^(11,14,35) Calcium and magnesium supplements are best taken with meals for this purpose, and to enhance their absorption.⁽³³⁾


A number of well-designed human intervention trials reveal that calcium supplementation...can lower blood pressure, particularly in sodium-sensitive hypertensive patients.

5. **Magnesium Supplementation** — Supplementation with 600 mg per day of magnesium has been shown to lower blood pressure in some, but not all, studies. Presently, a greater body of evidence exists for calcium supplementation than for magnesium. However, there is no risk in including 600 mg of magnesium in the management of hypertension (unless severe kidney disease is present).⁽¹⁵⁾



6. **Omega-3 Fat Supplementation** — Over sixty double-blind studies have demonstrated that either fish oil or flaxseed oil supplementation can be effective in lowering blood pressure. One tablespoon per day of flaxseed oil can lower systolic and diastolic blood pressure by up to 9 mm Hg.⁽¹⁶⁾ I generally recommend 1,000 mg of flaxseed oil (in capsule form), twice per day with meals.

7. **Garlic Extract Supplementation** — Supplementation with a garlic extract product that yields 4,000 mcg of allicin (1/2 to 1 clove of garlic) may help to lower blood pressure. Reductions of 20-30 mm Hg systolic and 10-20 mm Hg diastolic pressure have been demonstrated. However, this effect varies greatly among hypertensive subjects.^(2,17)



8. **Coenzyme Q10 Supplementation** — In recent years, a number of randomized, double-blind trials have demonstrated that Coenzyme Q10 (CoQ10) supplementation can effectively and consistently lower blood pressure in hypertensive subjects. CoQ10 is directly involved in the bioenergetic pathways of ATP production in heart muscle (myocardium). Research reveals that 39 percent of patients with high blood pressure have a deficiency of CoQ10. Supplementation with CoQ10 appears to correct this deficiency, correcting the underlying metabolic abnormality that leads to high blood pressure development. Most experts in this field believe that CoQ10 is able to lower blood pressure through its favourable influence on heart bioenergetic mechanisms and possibly relaxing vascular smooth muscle. Because CoQ10 corrects an

...Coenzyme Q10 supplementation can effectively and consistently lower blood pressure in hypertensive subjects...CoQ10 is able to lower blood pressure through its favorable influence on heart bioenergetic mechanisms and possibly relaxing vascular smooth muscle.

underlying metabolic defect that leads to high blood pressure, lowering of blood pressure usually requires 4 to 12 weeks of CoQ10 supplementation.^(18,19,20,21)

In a recent randomized, double blind trial among patients receiving antihypertensive medications, the addition of [60 mg of CoQ10, twice daily](#) was shown to markedly reduce both systolic and diastolic blood pressure. CoQ10 supplementation also reduced other risk factors for cardiovascular disease including lowering of fasting and 2-hr. plasma insulin, glucose, triglycerides, lipid peroxides and blood levels of malondialdehyde – a marker of free radical damage. The authors of the study conclude that CoQ10 decreases blood pressure possibly by decreasing oxidative stress (free radical generation) and insulin response in patients with known hypertension receiving conventional antihypertensive drugs. This study and others provide evidence that CoQ10 can be taken safely in conjunction with antihypertensive drugs to produce better blood pressure lowering outcomes, if necessary. ^(22,23,24)

In a recent randomized, double blind trial among patients receiving antihypertensive medications, the addition of 60 mg of CoQ10, twice daily, was shown to markedly reduce both systolic and diastolic blood pressure.

The [daily dosage of CoQ10](#) to aid in blood pressure lowering is usually 60 mg, twice per day, ⁽²²⁾ although 100 mg once per day has been tested ⁽¹⁶⁾ and in mild cases of hypertension 30-75 mg, once per day may be sufficient to normalize blood pressure.

9. [Hawthorn Extract Supplementation](#) — The hawthorn plant and its berries are a rich source of a unique strand of bioflavonoids, known as procyanidins. Like CoQ10, these procyanidins have been shown to reverse congestive heart failure by enhancing bioenergetic pathways within the heart muscle (myocardium). More recently, we have seen a number of intervention trials that demonstrate that [hawthorn extract supplementation](#) can also effectively reduce high blood pressure. The procyanidins in hawthorn act as cardiac glycoside agents that increase cyclic AMP and produce a vasodilation effect on arteries. The daily dosage required to lower blood pressure ranges from 100-250 mg, up to three times daily if taken as a sole antihypertensive agent. To ensure sufficient levels of its active constituents (procyanidins), the product must be standardized to 5 percent flavonoid content. (1-2% vitexin content) Usually two to four weeks is required in order to see a significant decline in blood pressure in hypertensive patients. ⁽²⁷⁾ Hawthorn is contra-indicated in patients taking digitalis or digoxin. ⁽³⁴⁾



Summary

The World Health Organization has promoted lifestyle modification as an effective method of reducing high blood pressure and overall cardiovascular risk. ⁽²⁴⁾ A summary of effective natural antihypertensive interventions include:

1. **Weight Loss** — Usually, only 10-15 lbs. of weight loss (in overweight subjects) will produce a significant blood pressure reduction in hypertensive patients.
2. **Salt Intake** — Limit to 2-3 gms per day
3. **Alcohol** — Limit alcohol consumption to less than two drinks per day and even less for women. (max. 9 drinks per week)
4. **Exercise** — Endurance exercise 30-60 minutes per session, a minimum of three times per week.
5. **Calcium Supplementation** — 1,000-1,500 mg per day (calcium carbonate or citrate), taken in divided doses of 500 mg per dose (with food).
6. **Magnesium Supplementation** — 600 mg per day (all at once or in divided doses, with food).
7. **Combination Flaxseed and Fish Oil** — 2,400-3,600 mg per day (two 1,200-mg capsules with meals).
8. **Coenzyme Q10** — 60 mg, twice per day is a popular treatment for hypertension.
9. **Hawthorn** — 75 mg, twice per day (std to 5% flavonoid content) can be used provided the patient is not also taking digitalis or digoxin.
10. **Garlic Extract Supplementation** (optional) — Endurance exercise 30-60 minutes per session, a minimum of three times per week.
11. **Fruits and Vegetables** — Increase fruit and vegetable intake to at least 5 servings per day.



The preceding recommendations can be used in conjunction with standard antihypertensive drugs if necessary. At present, there is sufficient evidence from well-designed medical trials to show that lifestyle interventions are successful in reducing or eliminating the need for pharmacologic therapy in a high percentage of hypertensive patients.

(29,30,31,32)

At present, there is sufficient evidence from well-designed medical trials to show that lifestyle interventions are successful in reducing or eliminating the need for pharmacologic therapy in a high percentage of hypertensive patients.

My suggestion is that you speak to your health practitioner about the appropriateness of these strategies in your individual case and seek his/her guidance as to how to access supplements that meet the requirements outlined in this review.

For more information on this or other related topics, visit Dr. Meschino's website at:

<http://www.meschinohealth.com/>

ADDITIONAL READINGS

(click on [http link below topic to view article](#))

1. **Coenzyme Q10 (CoQ10): An Essential Supplement for Cardiovascular Health After Forty**
http://www.meschinohealth.com/ArticleDirectory/CoEnzyme_Q10_%28CoQ10%29_An_Essential_Supplement_For_Cardiovascular_Health_After_Forty
2. **Reducing High Blood Pressure: Evidence-Based Natural Therapies**
http://www.meschinohealth.com/ArticleDirectory/Reducing_High_Blood_Pressure_Evidence-Based_Natural_Therapies
3. **Reishi Mushroom Extract and Immune Support**
http://www.meschinohealth.com/ArticleDirectory/Reishi_Mushroom_Extract_And_Immune_Support
4. **Hawthorn: Ancient Heart Disease Remedy Proven to Work in Recent Scientific Studies**
http://www.meschinohealth.com/ArticleDirectory/Hawthorn_Ancient_heart_disease_remedy
5. **How to Reduce High Blood Pressure Using Tested Natural Remedies**
http://www.meschinohealth.com/ArticleDirectory/How_To_Reduce_High_Blood_Pressure

- A Comprehensive Guide to Minerals
- A Comprehensive Guide to Herbs
- A Comprehensive Guide to Accessory Nutrients and Essential

References:

1. Quick Reference to Clinical Nutrition, Halpern, S. (ed.); Nutrition and Cardiovascular Disease; J.B. Lippincott Company, Philadelphia, 1987: 139-153
2. Canadian Guidelines for Cardiac Rehabilitation and Cardiovascular Disease Prevention (Canadian Assoc. of Cardiac Rehab.) 1st edition, 1999; 94-104
3. Fowler, F.E. Myocardial infarction in the 1990's; Postgraduate Medicine, May 1995; 97, 5: 135-146
4. Complete Guide to Prescription and Non-Prescription Drugs (1999 edition) Griffith H.W. The Body Press, 1998: 168-169, 194-195, 54-55
5. Murray, C.J.L.M., et al. Evidence-based health policy – lessons from the global burden of disease study. Science 1996; 274: 740-743
6. Joffres, M.R., et al. Awareness, treatment, and control of hypertension in Canada. Am J Hypertens. 1997; 10, (Pt-1): 1097-1102
7. 2000 Canadian hypertension recommendations (summary of recommendations affecting family physicians) – the Canadian Hypertension Recommendations Working Group. Canadian Family Physician. April 2001; 47: 793-794
8. Modern Nutrition in Health and Disease (sixth edition) Goodhart, R., and Shils, M. Lea and Febiger: 733
9. McCarron, D., et al. Body weight and blood pressure regulation. Am J Clin Nutr. 1996; 63 (suppl): 423-425
10. Pate, R.R., et al. Physical Activity and Public Health. JAMA. Feb. 1, 1995; 272, 5: 402-407
11. McCarron, D. Role of adequate dietary calcium intake in the prevention and management of salt-sensitive hypertension. Am J Clin Nutr. 1997; 62: 2 (suppl): 712-716
12. Cappuccio, F., et al. Double-blind randomized trial of modest salt restriction in older people. Lancet, 1997; 350; 9081: 850-854.
13. Graudal, N. et al. Effects of sodium restriction on blood pressure, rennin, aldosterone, catecholamines, cholesterol, and triglycerides. JAMA, 1998; 279: 1383-1391
14. Meese, R.B., et al. The inconsistent effects of calcium supplements upon blood pressure in primary hypertension. Am J Med Sci. 1987; 29: 4219-4224
15. Motoyama, T., et al. Oral magnesium supplementation in patients with essential hypertension. Hypertension, 1989; 13: 227-232
16. Murray, M., and Pizzorno, J. Encyclopedia of Natural Medicine (2nd edit.) Prima Publishing, 1997; 425-535
17. Foushee, D.B., et al. Garlic as a natural agent for the treatment of hypertension. A preliminary report. Cytobios. 1982; 34: 145-162
18. Digiesi, V., et al. Mechanism of action of Coenzyme Q10 in essential hypertension. Curr Ther Res. 1992; Res 51: 668-672
19. Langsjoen, P., et al. Treatment of essential hypertension with Coenzyme Q10. Mol Aspects Med. 1994; Med 15 (suppl): 265-272
20. Digiesi, V., et al. Coenzyme Q10 in essential hypertension. Mol Aspects Med. 1994; Med 15 (suppl): 257-263
21. McCarty, M.F. Coenzyme Q versus hypertension: does CoQ decrease endothelial superoxide generation? Med Hypotheses. 1999; 53, 4: 300-304
22. Singh, R.B., et al. Effect of hydrosoluble Coenzyme Q10 on blood pressure and insulin resistance in hypertensive patients with coronary artery disease. J Hum Hypertens. 1999; 13, 3: 203-208
23. Yamagami, T., et al. Bioenergetics in Clinical Medicine. Studies on Coenzyme Q10 and Essential Hypertension. Research Comm. in Chem. Path and Pharmacol 1975; 11, 2: 273-288
24. Yamagami, T., et al. Bioenergetics in Clinical Medicine, VIII. Administration of Coenzyme Q10 to patients with essential hypertension. Research Comm in Chem Path and Pharmacol. 1976; 14, 4: 721-727
25. Encyclopedia of Nutritional Supplements. Murray, M., PRIMA publishing, 1996: 300-301
26. Nutritional Influences on Illness. Werbach, M.R. Third Line Press., Inc. 1987: 227-240
27. Encyclopedia of Natural Medicine (2nd edit) Murray, M. and Pizzorno, J. Prima Publishing 1997: 524-535
28. Petrella, R.J. Lifestyle approaches to managing high blood pressure. Can Family Phys. 1999; 45: 1750-1755
29. Elmer, J.P., et al. Lifestyle intervention: results of the Treatment of Mild Hypertension Study. (TOHMS). Prev Med 1995; 24: 378-388
30. Stamler, R., et al. Nutritional therapy for high blood pressure. Final report of a four-year randomized controlled trial – the hypertension control program. JAMA. 1987; 257: 1484-1491
31. Iso, H., et al. Community-based education classes for hypertension control: a 1.5-year randomized controlled trial. Hypertension. 1996; 27: 968-974
32. Appel, L.J., et al. A clinical trial of the effects of dietary patterns on blood pressure (DASH-study) N Engl J Med 1997; 336: 1117-1124
33. Levenson, D., et al. A review of calcium preparations. Nutr Reviews. 1994; 52, 7: 221-232
34. Shariff, S., et al. Herbal Fervor and Vitamin Vigor: Herbs and vitamins for cardiac disease. Perspective in Cardiology. 2000; 16, 1: 21-29
35. McCarron, D., et al. Blood pressure response to oral calcium in persons with mild to moderate hypertension. A randomized, double-blind, placebo-controlled, crossover trial. Ann Intern Med, 1985; 103, 6: 825-831

Lower Your Blood Pressure Naturally

Vital Steps ANYONE Can Follow

Copyright © 2011

Dr. James Meschino, DC, MS, ND

All Rights Reserved

This eBook may not be reproduced in any form, in whole or in part,
without the consent of the publisher.

www.meschinohealth.com

1 - 888 - 494-1010

- A Comprehensive Guide to Minerals
- A Comprehensive Guide to Herbs
- A Comprehensive Guide to Accessory Nutrients and Essential



facebook.com/Meschino.Health



twitter.com/#!/DrJamesMeschino